



EXPAREL is available in 2 doses:
266 mg (20 mL) or 133 mg (10 mL)

EXPAREL Resources to Support Your Care Team and Patients

Pacira BioSciences, Inc. has created a variety of supporting resources with you and your EXPAREL patients in mind. Learn more about these complimentary items and how they can streamline your team and patient experience.

WANT TO ORDER COMPLIMENTARY ITEMS?

Contact your Pacira representative or scan the QR code with your mobile device.



EXPAREL Chart Sticker

Description:

A sticker affixed to the patient chart

For use by:



Nurses



Billers

- Nursing staff:** Local anesthetic utilization window reminder for multimodal therapy, PACU, and floor nurses
- Billers:** Procedure date tracking

ASC C9290 OR Baggie

Description:

Stores vials, wristband, and ASC OR log sticker



For use by:



Nurses



Billers

- Nursing staff:** Items to ready the OR and patient for the procedure, conveniently stored together
- Billers:** ASC OR log stickers to support billing documentation requirements

Patient Wristband

Description:

Wristband with EXPAREL administration timing

For use by:



Nurses



Patients

- Nursing staff:** Administration reminder
- Nursing staff:** Verification indicating when a patient may receive additional local anesthetic as part of an opioid-minimizing protocol
- Patients:** Safety in administration of additional local anesthesia as part of an opioid-sparing recovery process

ASC=ambulatory surgery center; HCPCS=Healthcare Common Procedure Coding System; OMFS=oral and maxillofacial surgery; OR=operating room; PACU=postanesthesia care unit.

Please see Important Safety Information on next page. Please [CLICK HERE](#) for full Prescribing Information.

OMFS Syringe Stickers

Description:

A sticker placed on syringes of EXPAREL ready for infiltration

For use by:



Nurses

- Nursing staff:** Time-stamped labels to distinguish syringes of EXPAREL being used in a procedure

ASC OR Log Stickers

Description:

A sticker for the OR log to document the amount of EXPAREL that is used AND unused in a procedure

For use by:



Nurses



Billers

- Nursing staff:** Used and unused dose to be provided by the surgeon
- Billers:** Both used and unused dose must be noted. **Billing is per mg, not per mL**
- Billers:** HCPCS code C9290 is used to bill for EXPAREL use in ASCs

Patient Discharge Instruction Tear Pad

Description:

Patient education resource communicating how to manage postsurgical pain after discharge

For use by:



Nurses



Patients

- Nursing staff:** Details the postsurgical care plan
- Patients:** Personalized take-home treatment and recovery information

Also available in Spanish

**NON-OPIOID
EXPAREL®**
(bupivacaine liposome injectable suspension)

NON-OPIOID
EXPAREL®
(bupivacaine liposome injectable suspension)

Indication

EXPAREL is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation.

Adverse reactions with an incidence greater than or equal to 10% following EXPAREL administration via infiltration in pediatric patients six to less than 17 years of age were nausea, vomiting, constipation, hypotension, anemia, muscle twitching, vision blurred, pruritis, and tachycardia.

If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine.

EXPAREL is not recommended to be used in the following patient populations: patients <6 years old for infiltration, patients younger than 18 years old for interscalene brachial plexus nerve block, and/or pregnant patients.

Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability, which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Methemoglobinemia: Cases of methemoglobinemia have been reported with local anesthetic use.

Full Prescribing Information is also available at www.EXPAREL.com.