



BioXclude®

SOCKET PRESERVATION

Comprehensive User Guidance for:

- Minimally Invasive Socket Preservation
- Socket Preservation with Flap Elevation
- Suturing & Post-Operative Instruction


SIMPLE EFFECTIVE VERSATILE



Socket Preservation Handling Guide

Step-by-step handling instructions for BioXclude®
Dehydrated human deepithelialized amnion-chorion membrane (ddACM)


Minimally Invasive Socket Preservation




1. A contained extraction socket should be filled with bone graft to the height of the crestal walls. BioXclude should be placed on top of bone graft material with minimal flap reflection.




2. Place dry, untrimmed 8x8 mm (anterior) or 12x12 mm (posterior) BioXclude using dry forceps. Orientation during placement does not matter. BioXclude may be placed UP or DOWN.




3. BioXclude will hydrate and adapt to bone particulate. Drops of irrigant (sterile saline) can speed up membrane hydration. Damp gauze can be used to help adapt BioXclude over the underlying graft while absorbing excess fluids.



4. It is easiest to not manipulate the membrane and instead suture from the inside of the socket (connective tissue side out) to avoid nicking the membrane first.



5. Using this reverse, inverted suturing method in a figure 8 (one for an anterior site and two for a posterior site) will approximate the tissues over the membrane. A PTFE suture is recommended. Continue to blot with damp gauze as needed.



6. Use a wetted instrument (Buser periosteal elevator works well) to tuck the edges of BioXclude as necessary. BioXclude only needs to be 1 mm under the gingival margin.

Toolkit

1. Adson Forcep
2. Minnesota Retractor
3. Molt #9 Periosteal Elevator
4. Curette
5. Buser Periosteal Elevator
6. Needle Holder
7. Scissor
8. Cotton Forcep
9. SS Medicine Cup (or dappen dish)
10. SS Iodine bowl (for saline)
11. Sterile Gauze
12. Suture
13. Monoject Syringe
14. Sterile Saline



Welcome to handling, **without rules:**


- » No need to trim, tack or suture
- » No orientation - place BioXclude “up” or down, fold it, or allow the membrane to “bunch” up
- » Safely touch tooth, root or implant surfaces
- » Place over or under other membranes or mesh
- » Self-adheres and adapts

Socket Preservation with **Flap Elevation**


Helpful Hint:

Bone particulate with a mineralized component is commonly used.


Note the necessity to overbulk the buccal defect with bone particulate due to the likelihood of resorption.



1. BioXclude is placed last, after all of the bone particulate is placed. There is no need to trim BioXclude - it is safe to touch adjacent tooth surfaces.



2. BioXclude is brought to the site dry. Choose a BioXclude size to extend over all graft material and onto native buccal bone, over the crest and tucked lingually.




3. Use an instrument to anchor BioXclude on the crest. A monoject syringe with sterile saline can be used to hydrate the membrane as needed.



4. BioXclude will naturally adapt and adhere to bone particulate and to adjacent native bone.



5. A damp gauze may be pressed against the site to absorb additional fluid to reapproximate the flap without disrupting the membrane.






6. After vertical releases are sutured, non-primary closure can be obtained using 4.0 PTFE suture, and a reverse figure eight technique (see ‘Suturing Guide’ instructions below).

Suturing Guide

Reverse or “inverted” sutures pull the tissue both inward and downward. In an open socket this is ideal for membrane retention.

This method also greatly decreases the potential for nicking the membrane.



Note: Each pass begins from inside the socket (connective tissue side).

BioXclude® Size Choices

CHOOSING THE RIGHT SIZE:




8x8 mm



12x12 mm




10x20 mm



15x20 mm



15x25 mm



20x30 mm

No Flap Elevation:

Tuck 1 mm under gingival margin

Flap Elevation:

Cover all graft material and extend onto native bone 3 mm

Post Operative Guidance


When the membrane is exposed to the oral environment:

- » **No** rinsing, swishing, spitting, or sucking through a straw for the **first 3 days**. These actions can dislodge the membrane.
- » **No chlorhexidine or OTC mouth rinses**. Oral rinses are used to kill bacteria. To varying degrees, oral rinses adversely impact the health of gingival cells, thus slowing wound closure. Fortunately, Purion® processed amnion-chorion allografts have demonstrated natural anti-microbial properties.
- » After **3 days**, gentle rinsing with tap water is recommended for the next **7 days**. Only tap water should be used during this time frame. After **10 days** post-operatively, the patient may begin using an oral rinse for plaque control.

Appearance During Healing


Using BioXclude with Non-Primary Closure

Variation in healing appearance, including translucent, opaque, and yellowish appearance, are all normal and common.



4 day post-op

Anthony Del Vecchio, DDS




3 day post-op

Dan Holtzclaw, DDS, MS




4 day post-op

Anthony Del Vecchio, DDS




4 day post-op

Dan Holtzclaw, DDS, MS




10 day post-op

Matthew J. Fien, DDS




2 week post op

Dean Licenblat, BDent, MSc



2 week post op*
(*pt is a smoker)

Vinay Bhide, DDS, MSc




10 day post-op

Vinay Bhide, DDS, MSc


Your Membrane Choice Matters

Superior bone growth was observed when comparing **BioXclude®** dehydrated human deepithelialized amnion-chorion membrane to **dPTFE** in socket preservation


New Osteoid/Bone Formation



Bone Volume Density



Patient Discomfort



The Keys to BioXclude's Success

Amnion-Chorion Membrane

- Immunoprivileged allograft tissue
- No history of graft rejection
- Natural tissue barrier
- Intact basement membranes
- Active growth factors and chemokines
- Known to play critical roles in tissue repair and regeneration
- Antibacterial properties

Purion® Processed

- The gold standard in placental tissue processing
- Safely and gently ensures that the key elements associated with healing are preserved
- Proprietary to BioXclude®

The Difference is in the Deepithelialization

- Exposes underlying basement membranes
- Improves cellular attachment
- Allows membrane to be placed "up" or "down" at the treatment site

Randomized Clinical Trials

- Samer, A. (2020). *Journal of Oral Implantology*
- Hassan, M. (2017). *International Journal of Oral and Maxillofacial Implants*

Case Series

- Maksoud, M.A (2018). *Clinical Advances in Periodontics*
- Holtzclaw, D. (2014). *Compendium of Continuing Education in Dentistry*
- Wallace, S. (2011). *Journal of Implant and Advanced Clinical Dentistry*
- Holtzclaw, D. (2011). *Journal of Implant and Advanced Clinical Dentistry*

Case Reports

- Cullum, D. et al. (2019). *Compendium of Continuing Education in Dentistry*
- Prakasam, S. (2017). *Decisions in Dentistry*
- Wallace, S. (2010). *Journal of Implant and Advanced Clinical Dentistry*



SIMPLE
EFFECTIVE
VERSATILE



9 Total Publications



on **socket preservation** ranging from case reports to randomized clinical trials


BioXclude® Socket Preservation: Clinical Cases




Matthew J. Fien, DDS
Fort Lauderdale, FL
@fienodontics

**John Kim,
DMD, MS, PA**
Rocky Mount, NC
@rockymountperio






Hard Tissue Histology



Surface area of new bone/host bone = 51.1%


Surface area of residual graft material = 24.2%

Surface area of connective tissue = 24.7%




4 mm

Soft Tissue Histology



This specimen represents the classic keratinized stratified squamous epithelium. There was little to no parakeratin and the stratum granulosum was not well developed; however, this presentation is still within normal limits. Parakeratin and a well developed granular cell layers are more typical of attached gingiva that receives "stimulation" from daily tooth brushing and mastication



2 week healing



2 week healing



Michael Block, DMD

CENTER FOR DENTAL RECONSTRUCTION | METAIRIE, LA

"I have been using BioXclude to cover my extraction site grafts and to cover larger grafts for ridge augmentation. **My incision dehiscence rate is almost zero.** The soft tissue healing using this material is very strong. **I recommend it without reservations.**"

Matthew Fien, DDS

FIENODONTICS | PLANTATION, FL



"**Bioxclude is an incredible biomaterial** and it has changed the way I practice. **The applications are endless**"



Dan Holtzclaw, DDS, MS

DENTAL IMPLANT CENTER | AUSTIN, TX

"**I have used it thousands and thousands of times** - I have used it on family members, I have used it on friends. **I know that it works.** Now, there are many, many, **published studies** that also show that this material works and there are **histological studies** to back it up."

John Alonge, DDS, MS

ORAL SURGERY OF ERIE | ERIE, PA



"**My patients have experienced far less pain**, swelling and bad taste with this membrane than with anything else I have used. I have noticed a **marked reduction in mucosa inflammation** and **faster healing.**"

