Section Calmoseptine® Ointment

MULTIPURPOSE MOISTURE BARRIER



Available in: 4 oz Tubes, 2.5 oz Tubes, 2.5 oz Jars, 20g Tubes and 1/8 oz Packets

Calmoseptine, Inc.

16602 Burke Lane • Huntington Beach, California 92647 • U.S.A.

(800) 800-3405 • FAX (714) 840-9810 www.calmoseptine.com

Stalmoseptine® Ointment

PROTECTS AND HELPS HEAL SKIN IRRITATIONS FROM:

- INCONTINENCE OF URINE OR FECES
- FEEDING TUBE SITE LEAKAGE

• WOUND DRAINAGE

FECAL OR VAGINAL FISTULAS

MINOR BURNS, SCRAPES

DIAPER RASH

MOISTURE, SUCH AS PERSPIRATION

REMARKABLE RESULTS IN JUST OVER 48 HOURS CASE STUDY #00305573 - FECAL INCONTINENCE



DAY 1

On day 1, this 81 year old female was admitted with multiple dermal ulcers, diarrhea (8-10 stools daily), and a Foley catheter for urinary incontinence, 8x10 inches of perianal skin was red and raw from fecal erosion. Treatment with Calmoseptine[®] Ointment was started.



DAY 3

By day 3, the patient was less agitated and more comfortable. A 1x3 inch reddened area around the anus was all that remained. Diarrhea continued, but with the use of Calmoseptine® Ointment applied twice daily, healing took place rapidly.

Although this patient's urinary incontinence was controlled by a catheter, she experienced intermittent diarrhea over the next 15 months. The use of Calmoseptine[®] Ointment prevented skin breakdown. - Case study by Odette Birth, RN, CETN

DIRECTIONS

- 1. Cleanse skin gently with mild skin cleanser. Pat dry or air dry.
- 2. Apply a thin layer of Calmoseptine® Ointment to reddened or irritated skin 2-4 times daily, or after each incontinent episode or diaper change to promote comfort, and long lasting protection. (If Calmoseptine® Ointment residue remains on irritated skin after gentle cleansing, it is recommended to leave the residue on the skin and apply an additional layer as needed until the skin is healed.) Calmoseptine® Ointment is easily removed with baby oil or mineral oil when necessary.

ORDERING INFORMATION		
NDC #	SIZE	PER CASE
0799-0001-02	2.5 oz. (71g) tube	12
0799-0001-03	2.5 oz. (71g) jar	12
0799-0001-04	4.0 oz. (113g) tube	12
0799-0001-05	1/8 oz. (3.5g) packe	t 144
0799-0001-06	0.7 oz. (20g) tube	12

ACTIVE INGREDIENTS:

Menthol 0.44% External analgesic / Anti-itch Zinc Oxide 20.6% Skin protectant / Anorectal astringent OTHER INGREDIENTS: calamine, chlorothymol, glycerin, lanolin, phenol, sodium bicarbonate and thymol in a suitable ointment base

ENCOURAGE BASIC CONCEPTS OF SKIN CARE

- Keep skin clean, moisturized and dry.
 Adequate "pressure relief" for bedridden and wheelchair bound patients.
- · Assessment and monitoring of nutritional status by a dietitian. · Adequate fluid intake (6-8 glasses of water daily).
- · Consultation by an ET Nurse (Enterostomal Therapist) or skin specialist.
- Protect yourself and the patient by observing universal blood and body fluid precautions.

WARNING: For external use only. Not for deep or puncture wounds. Avoid contact with eyes. Keep out of reach of children. In case of accidental ingestion contact a physician or poison control center immediately. If condition worsens or does not improve within 7 days, consult a doctor.

Calmoseptine, Inc. 16602 Burke Lane • Huntington Beach, California 92647 • U.S.A. US -21 (800) 800-3405 · FAX (714) 840-9810 www.calmoseptine.com